

NOVARTIS AG
Form 6-K
June 06, 2011

SECURITIES AND EXCHANGE COMMISSION

Washington, D.C. 20549

FORM 6-K

**REPORT OF FOREIGN PRIVATE ISSUER
PURSUANT TO RULE 13a-16 or 15d-16 OF
THE SECURITIES EXCHANGE ACT OF 1934**

Report on Form 6-K dated June 5, 2011

(Commission File No. 1-15024)

Novartis AG

(Name of Registrant)

Lichtstrasse 35

4056 Basel

Switzerland

(Address of Principal Executive Offices)

Indicate by check mark whether the registrant files or will file annual reports under cover of Form 20-F or Form 40-F:

Edgar Filing: NOVARTIS AG - Form 6-K

Form 20-F: **Form 40-F:**

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(1):

Yes: **No:**

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(7):

Yes: **No:**

Indicate by check mark whether the registrant by furnishing the information contained in this form is also thereby furnishing the information to the Commission pursuant to Rule 12g3-2(b) under the Securities Exchange Act of 1934.

Yes: **No:**

Novartis International AG

Novartis Global Communications
CH-4002 Basel
Switzerland
<http://www.novartis.com>

- Investor Relations Release -

Novartis drug Glivec® shows significant overall survival benefit for patients with GIST after three years of adjuvant treatment vs. one year

- *Data show 66% recurrence-free survival and 92% overall survival at five years following three years of adjuvant therapy with Glivec in patients with resected KIT+ GIST(1)*
- *First large Phase III clinical trial to demonstrate survival benefits of extended treatment with Glivec for three years compared to one year following surgery*
- *Important finding for treatment of GIST patients who are at risk of recurrence following complete resection of primary tumor*

Basel, June 5, 2011 Novartis announced new data today showing a significant improvement in both recurrence-free survival and overall survival for patients taking Glivec® (imatinib)* for three years after surgery to remove KIT (CD117)-positive gastrointestinal stromal tumors (GIST) compared to one year of treatment. These data will be presented at the 47th American Society of Clinical Oncology (ASCO) plenary session on Sunday, June 5.

The results show that at five years 66% of patients taking Glivec for three years remained free of cancer recurrence (primary endpoint) compared to 48% who had received Glivec for only one year ($p < .0001$). Moreover, 92% of patients taking Glivec for three years were alive (secondary endpoint) compared to 82% who had received Glivec for only one year ($p = .019$). Median patient follow-up was 54 months.(1)

The 400-patient Phase III trial, conducted by the Scandinavian Sarcoma Group (SSG) and the Sarcoma Group of the Arbeitsgemeinschaft Internistische Onkologie (AIO), is the first prospective multicenter clinical trial to demonstrate a survival benefit of adjuvant Glivec therapy for KIT+ GIST with extended three years of therapy relative to one year of therapy. The side effect profile in the clinical trial was consistent with that reported in previous studies with Glivec.

This study confirms the hypothesis that extending the duration of Glivec treatment for patients following surgery improves recurrence-free survival. For the first time, an effect on overall survival was found, said Heikki Joensuu, M.D., Ph.D., Professor, Oncology, University of Helsinki and principal investigator of the study. Results from this trial may positively impact clinical practice by helping physicians create the

optimal treatment plan for patients with operable KIT+ GIST.

Gastrointestinal stromal tumors are a rare, life-threatening cancer of the gastrointestinal tract. The major cause of GIST is an abnormal form of the protein KIT,(2) which causes cells to grow uncontrollably and become cancerous. Patients with GIST are at risk of recurrence following complete resection of primary GIST.(3)

Over the past nine years Glivec has provided KIT+ GIST patients with the first effective drug treatment option in the metastatic setting and later in the adjuvant setting, said Hervé Hoppenot,

President, Novartis Oncology. Now we see exciting new data showing that by extending post-surgical treatment duration to three years, Glivec has significant impact on overall survival in patients with KIT+ GIST. This is important news for GIST patients and the GIST community.

Study details

The SSG XVIII clinical trial was conducted by the Scandinavian Sarcoma Group (SSG), and the Sarcoma Group of the Arbeitsgemeinschaft Internistische Onkologie (AIO). SSG XVIII is a multicenter, prospective, randomized study for the evaluation of adjuvant treatment with Glivec of histologically verified KIT+ GIST with a greater than 50% risk of GIST recurrence despite complete removal of all macroscopic GIST tissue at surgery.(4)

The primary endpoint of the study was to compare the recurrence-free survival in GIST patients with a greater than 50% estimated risk of disease recurrence within the first five years following the diagnosis and treatment with adjuvant Glivec for 12 or 36 months. The secondary endpoints included overall survival and treatment safety.(1)

Four hundred patients entered the study and the median follow-up was 54 months. Recurrence-free survival was longer in the 36-month group compared to the 12-month group (HR 0.46, 95% CI 0.32-0.65; $p < .0001$; five-year recurrence-free survival 66% vs. 48%, respectively). Patients assigned to 36 months of Glivec had longer overall survival (HR 0.45, 95% CI 0.22-0.89; $p = .019$; five-year overall survival 92% vs. 82%, respectively). Glivec was generally well tolerated. The proportion of patients who discontinued Glivec during the assigned treatment period for reasons other than GIST recurrence was 26% in the 36-month group and 13% in the 12-month group.(1)

About Glivec

Glivec is approved in more than 110 countries, including the US, EU and Japan, for the treatment of all phases of Ph+ CML. Glivec is also approved in the US, EU and other countries for the treatment of patients with KIT (CD117)-positive gastrointestinal stromal tumors (GIST), which cannot be surgically removed and/or have already spread to other parts of the body (metastasized). In the US and EU, Glivec is approved for the post-surgery treatment of adult patients following complete surgical removal of KIT (CD117)-positive gastrointestinal stromal tumors.

Not all indications are available in every country and indications may differ.

Glivec Important Safety Information

Glivec can cause fetal harm when administered to a pregnant woman. Women should not become pregnant, and should be advised of the potential risk to the unborn child.

Edgar Filing: NOVARTIS AG - Form 6-K

Glivec is often associated with edema (swelling) and serious fluid retention. Studies have shown that edema (swelling) tended to occur more often among patients who are 65 and older or those taking higher doses of Glivec.

Cytopenias (reduction or lack of certain cell elements in blood circulation), such as anemia, have occurred. If the cytopenia is severe, your doctor may reduce your dose or temporarily stop your treatment with Glivec.

Severe congestive heart failure and left ventricle dysfunction have been reported, particularly in patients with other health issues and risk factors. Patients with heart disease or risk factors will be monitored and treated for the condition.

Severe liver problems (hepatotoxicity) may occur. Cases of fatal liver failure and severe liver injury requiring liver transplants have been reported with both short-term and long-term use of Glivec.

Bleeding may occur. Severe gastrointestinal (GI) bleeding has been reported in patients with KIT+ GIST. GI tumor sites may be the cause of this bleeding.

In patients with hypereosinophilic syndrome (a condition with increased eosinophils, which are a type of white blood cell) and heart involvement, cases of heart disease (cardiogenic shock/left ventricular dysfunction) have been associated with the initiation of Glivec therapy.

Skin reactions, such as fluid-filled blisters, have been reported with the use of Glivec. Clinical cases of hypothyroidism (reduction in thyroid hormones) have been reported in patients taking levothyroxine replacement with Glivec.

Long-term use may result in potential liver, kidney, and/or heart toxicities; immune system suppression may also result from long-term use.

GI perforation (small holes or tears in the walls of the stomach or intestine), in some cases fatal, has been reported.

Growth retardation has been reported in children taking Glivec. The long-term effects of extended treatment with Glivec on growth in children are unknown.

Cases of tumor lysis syndrome (TLS), which refers to a metabolic and electrolyte disturbance caused by the breakdown of tumor cells, have been reported and can be life-threatening in some cases.

Almost all patients with KIT+ GIST treated with Glivec experience side effects at some time. Some common side effects you may experience are fluid retention, muscle cramps or pain and bone pain, abdominal pain, loss of appetite, vomiting, diarrhea, decreased hemoglobin, abnormal bleeding, nausea, fatigue and rash.

Glivec is sometimes associated with stomach or intestinal irritation. Glivec should be taken with food and a large glass of water to minimize this problem. There have been rare reports, including deaths, of stomach or intestinal perforation (a small hole or tear).

If you are experiencing any of the mentioned side effects, please be sure to speak with your doctor immediately.

Do not take any other medications without talking to your doctor or pharmacist first, including Tylenol® (acetaminophen); herbal products (St. John's wort, Hypericum perforatum); Coumadin® (warfarin sodium); rifampin; erythromycin; metoprolol; ketoconazole; and Dilantin® (phenytoin). Taking these with Glivec may affect how they work, or affect how Glivec works.

Edgar Filing: NOVARTIS AG - Form 6-K

You should also tell your doctor if you are taking or plan to take iron supplements. Patients should also avoid grapefruit juice and other foods that may affect how Glivec works.

Please see full Prescribing Information.

Disclaimer

The foregoing release contains forward-looking statements that can be identified by terminology such as will, may, or similar expressions, or by express or implied discussions regarding potential new indications or labeling for Glivec or regarding potential future revenues from Glivec. You should not place undue reliance on these statements. Such forward-looking statements reflect the current views of management regarding future events, and involve known and unknown risks, uncertainties and other factors that may cause actual results with Glivec to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that Glivec will be submitted or approved for any additional indications or labeling in any market. Nor can there be any guarantee that Glivec will

achieve any particular levels of revenue in the future. In particular, management's expectations regarding Glivec could be affected by, among other things, unexpected regulatory actions or delays or government regulation generally; unexpected clinical trial results, including unexpected new clinical data and unexpected additional analysis of existing clinical data; the company's ability to obtain or maintain patent or other proprietary intellectual property protection; competition in general; government, industry and general public pricing pressures; the impact that the foregoing factors could have on the values attributed to the Novartis Group's assets and liabilities as recorded in the Group's consolidated balance sheet, and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those anticipated, believed, estimated or expected. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

About Novartis

Novartis provides healthcare solutions that address the evolving needs of patients and societies. Focused solely on healthcare, Novartis offers a diversified portfolio to best meet these needs: innovative medicines, eye care, cost-saving generic pharmaceuticals, consumer health products, preventive vaccines and diagnostic tools. Novartis is the only company with leading positions in these areas. In 2010, the Group's continuing operations achieved net sales of USD 50.6 billion, while approximately USD 9.1 billion (USD 8.1 billion excluding impairment and amortization charges) was invested in R&D throughout the Group. Headquartered in Basel, Switzerland, Novartis Group companies employ approximately 119,000 full-time-equivalent associates and operate in more than 140 countries around the world. For more information, please visit <http://www.novartis.com>.

References

- (1) Joensuu H, et al. Twelve vs. 36 months of adjuvant imatinib (IM) as treatment of operable GIST with a high risk of recurrence: Final results of a randomized trial (SSGXVIII/AIO). 47th Annual Meeting of the American Society of Clinical Oncology. Abstract No. LBA1. June 5, 2011.
- (2) Gomes AL, Bardales RH, Milanezi F, Reis RM, Schmitt F. Molecular analysis of c-KIT and PDGFRA in GISTs diagnosed by EUS. *Am J Clin Pathol.* 2007 Jan;127(1):89-96.
- (3) DeMatteo RP, Lewis JJ, Leung D, Mudan SS, Woodruff JM, Brennan MF. Two hundred gastrointestinal stromal tumors: recurrence patterns and prognostic factors for survival. *Ann Surg.* 2000 Jan;231(1):51-8.
- (4) Study Comparing 12 Months Versus 36 Months of Imatinib in the Treatment of Gastrointestinal Stromal Tumor (GIST). Available at <http://clinicaltrials.gov/show/NCT00116935>. Accessed on April 11, 2011.

Novartis is on Twitter. Sign up to follow @Novartis at <http://twitter.com/novartis>.

###

Novartis Media Relations

Central media line : +41 61 324 2200

Eric Althoff

Novartis Global Media Relations

+41 61 324 7999 (direct)

+41 79 593 4202 (mobile)

eric.althoff@novartis.com

Sabrina Oei

Novartis Oncology

+1 862.778.6387 (direct)

+1 862.210.0993 (mobile)

sabrina.oei@novartis.com

e-mail: media.relations@novartis.com

For Novartis multimedia content, please visit www.thenewsmarket.com/Novartis

For questions about the site or required registration, please contact: journalisthelp@thenewsmarket.com.

Novartis Investor Relations

Central phone:

Susanne Schaffert

Pierre-Michel Bringer

Thomas Hungerbuehler

Isabella Zinck

+41 61 324 7944

+41 61 324 7944

+41 61 324 1065

+41 61 324 8425

+41 61 324 7188

North America:

Richard Jarvis

Jill Pozarek

Edwin Valeriano

+1 212 830 2433

+1 212 830 2445

+1 212 830 2456

e-mail: investor.relations@novartis.com

e-mail: investor.relations@novartis.com

SIGNATURES

Pursuant to the requirements of the Securities Exchange Act of 1934, the registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

Novartis AG

Date: June 5, 2011

By:

/s/ MALCOLM B. CHEETHAM

Name:

Malcolm B. Cheetham

Title:

Head Group Financial Reporting and Accounting